



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 9/00	A2	(11) International Publication Number: WO 00/06706 (43) International Publication Date: 10 February 2000 (10.02.00)
(21) International Application Number: PCT/EP99/05413 (22) International Filing Date: 28 July 1999 (28.07.99) (30) Priority Data: 9816639.0 30 July 1998 (30.07.98) GB (71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). (72) Inventors; and (75) Inventors/Applicants (for US only): REVENKOVA, Ekaterina [RU/CH]; Peter Rot-Strasse 76, CH-4058 Basel (CH). PASZOWSKI, Jerzy [PL/CH]; Blauenweg 10, CH-4224 Nenzlingen (CH). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: MAP KINASE PHOSPHATASE MUTANT (57) Abstract <p>The present invention relates to DNA encoding proteins contributing to the regulation of a plant's response to DNA damage. DNA according to the invention comprises an open reading frame encoding a protein characterized by a stretch of amino acids or component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO:3. Preferably the DNA encodes a MAP kinase phosphatase.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

MAP kinase phosphatase mutant

The present invention relates to DNA encoding proteins contributing to the regulation of a plant's response to abiotic stress and in particular genotoxic stress.

Cells of all organisms have evolved a series of DNA repair pathways which counteract the deleterious effects of DNA damage and are triggered by intricate signal cascades. To be able to modify or improve DNA repair using gene technology it is necessary to identify key proteins involved in said pathways or cascades. Therefore it is the main object of the present invention to provide DNA comprising an open reading frame encoding such a key protein.

DNA according to the present invention comprises an open reading frame encoding a protein characterized by a stretch of amino acids or component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO: 3. The protein characterized by SEQ ID NO: 3 is tracked down with the help of a T-DNA tagged Arabidopsis mutant showing hypersensitivity to methyl methanesulfonate (MMS). Said hypersensitivity as well as an observed hypersensitivity to other DNA damaging treatments such as UV light is indicative of the proteins' involvement in the repair of DNA damage, or in signaling pathways implicated in the response to similar genotoxic stress. The mutant is also sensitive to elevated temperature and anti-oxidant N-acetylcysteine. The mutant is not sensitive to osmotic shock, increased salinity, oxidative stress or elevated ethylene levels. An important characteristic of the mutant is cell death in response to growth in small closed vessels. This phenotype can be complemented by addition of abscisic acid (ABA) to the growth media. Furthermore, the mutant is more sensitive to exogenously applied ABA compared with the wild type which supports the notion that the genes disclosed by the present invention (SEQ ID NO: 1) are involved in stress signaling mediated by ABA.

Sequence alignments of SEQ ID NO: 3 using commercially available computer programs such as BLASTP of the NCBI BLAST family of programs or TFASTA or BestFit of the Wisconsin Package Software, all based on well known algorithms for sequence identity or similarity searches, reveal that stretches of SEQ ID NO: 3 (component sequences) having more than 100 and preferably between 120 to 250 amino acids length can show between

20% and almost 40% sequence identity to aligned stretches of known phosphatases, particularly phosphotyrosine phosphatases, MAP kinase phosphatases or dual specificity phosphatases. Protein phosphatases are classified by their substrate specificities as either phosphoserine/threonine phosphatases (PSTPs) or phosphotyrosine phosphatases (PTPs). The dual specificity phosphatases (DSPs) dephosphorylate both phosphotyrosine and phosphoserine/threonine residues and represent a subfamily of PTPs. MAP kinase phosphatases (MKPs) belong to the family of DSPs. The sequence VHCCQGVSR (SEQ ID NO: 4) found in SEQ ID NO: 3 can be interpreted as corresponding to the mammalian sequence motif IHGXGXXRS (SEQ ID NO: 5) defining the family of PTPs, wherein the Ile at the first position can be replaced by Val and the Ser at the last position can be replaced by Thr

The present invention defines a new protein family the members of which are characterized by component amino acid sequences of more than 100 amino acid length showing 40% or higher amino acid sequence identity to aligned component sequences of SEQ ID NO: 3. Preferably said component sequences are of more than 120, more than 160 or even more than 200 amino acids length. The amino acid sequence identity is preferably higher than 50% or even higher than 55%. Most preferred are identities higher than 70%.

An example of DNA according to the present invention is described in SEQ ID NO: 1. The amino acid sequence of the protein encoded is identical to SEQ ID NO: 3. After alignment a stretch of the protein having about 140 amino acids shows 36% sequence identity to the MKP-1 protein described by Sun et al (Cell 75: 487-493, 1993). The identity determined after alignment with MKP-2 and MKP-3 is determined as 34% and 26%, respectively. Thus, according to the present invention a protein family related to MAP kinase phosphatases can be defined the members of which after alignment of a stretch of more than 100 amino acids length show 40% or higher amino acid sequence identity to SEQ ID NO: 3. Preferably, the amino acid sequence identity is higher than 50% or even higher than 55%. When making multiple sequence alignments, certain algorithms can take into account sequence similarities, such as same net charge or comparable hydrophobicity/hydrophilicity of the individual amino acids, in addition to sequence identities. The resulting values of sequence similarities, as compared to sequence identities, can help to assign a protein to the correct protein family in border-line cases. Proteins of particular interest, within the scope of the present invention, are MAP kinase phosphatases the amino acid sequence of which comprises at least one of the following characteristic amino acid subsequences:

- (a) TSILYDVFDYFEDV (SEQ ID NO: 6)
- (b) FVHCCQGVSRST (SEQ ID NO: 7)
- (c) FVHC (SEQ ID NO: 8)
- (d) QGVSRST (SEQ ID NO: 9)
- (e) YFKSD (SEQ ID NO: 10)

DNA encoding proteins belonging to the new protein family according to the present invention can be isolated from monocotyledonous and dicotyledonous plants. Preferred sources are corn, sugar beet, sunflower, winter oilseed rape, soybean, cotton, wheat, rice, potato, broccoli, cauliflower, cabbage, cucumber, sweet corn, daikon, garden beans, lettuce, melon, pepper, squash, tomato, or watermelon. However, they can also be isolated from mammalian sources such as mouse or human tissues. The following general method, can be used, which the person skilled in the art will normally adapt to his specific task. A single stranded fragment of SEQ ID NO: 1 or SEQ ID NO: 2 consisting of at least 15, preferably 20 to 30 or even more than 100 consecutive nucleotides is used as a probe to screen a DNA library for clones hybridizing to said fragment. The factors to be observed for hybridization are described in Sambrook et al, Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press, chapters 9.47-9.57 and 11.45-11.49, 1989. Hybridizing clones are sequenced and DNA of clones comprising a complete coding region encoding a protein with more than 40% sequence identity to SEQ ID NO: 3 is purified. Said DNA can then be further processed by a number of routine recombinant DNA techniques such as restriction enzyme digestion, ligation, or polymerase chain reaction analysis.

The disclosure of SEQ ID NO: 1 enables a person skilled in the art to design oligonucleotides for polymerase chain reactions which attempt to amplify DNA fragments from templates comprising a sequence of nucleotides characterized by any continuous sequence of 15 and preferably 20 to 30 or more basepairs in SEQ ID NO: 1. Said nucleotides comprise a sequence of nucleotides which represents 15 and preferably 20 to 30 or more basepairs of SEQ ID NO: 1. Polymerase chain reactions performed using at least one such oligonucleotide and their amplification products constitute another embodiment of the present invention.

Knowing the nucleotide sequence of the *Arabidopsis* MKP1 gene and the amino acid sequence of the encoded protein it is possible to identify proteins interacting with AtMKP1 and to clone their corresponding genes using well known techniques. For example radioactively labeled AtMKP1 protein can be used for interactive cloning on cDNA expression libraries. AtMKP1 protein or parts thereof can be used to generate polyclonal or monoclonal antibodies specific for AtMKP1. The *AtMKP1* gene can be used to generate variants of AtMKP1 protein tagged with GST, MYK or His. Said antibodies and MKP1 variants allow to isolate native protein complexes by immunoprecipitation and to determine sequences of proteins present in these complexes by micro-sequencing. The resulting sequence information can in turn be used to clone corresponding genes. Alternatively, said antibodies or tagged MKP1 variants can be used to screen epitope libraries for epitopes which interact with AtMKP1 protein. The AtMKP1 protein and parts thereof, in particular the N-terminal 490 amino acid region and the C-terminal 492 amino acid region can also be used to search for interacting proteins with a Two-hybrid system (e.g. in yeast, in mammalian cells, or in bacteria). This allows to obtain sequence information about interacting proteins.

Based on the disclosed finding that AtMKP1 proteins are involved in a plant's abiotic environmental stress response, it becomes possible to engineer the corresponding signaling pathway, of which AtMKP1 is a part, to be chemically regulated due to chemical activation or repression of transgenes encoding AtMKP1 or proteins interacting therewith. Such plants can be obtained by transformation with the corresponding genes under control of chemically inducible promoters. Application of inducers is expected to modify the activity of the AtMKP1 signaling pathway and to result in altered adaptation to abiotic environmental stress. Alternatively, AtMKP1 protein or its interacting proteins can be used as targets for chemicals inhibiting or stimulating their activities which again is expected to modify abiotic stress responses.

EXAMPLES:

Example 1: *Cloning of the gene responsible for the mkp1 mutant phenotype*

Arabidopsis T-DNA insertion lines as produced by the INRA-Versailles and available from the Nottingham Arabidopsis Stock Center (NASC) are screened for sensitivity to methyl methanesulfonate (MMS) at a concentration of 100 ppm as described by Masson et al,

Genetics 146: 401-407, 1997. Plants which die in the presence of 100 ppm MMS are found in the family AAN4. Thus, the corresponding T-DNA insertion mutation is assumed to give rise to this hypersensitive phenotype. This assumption is supported by genetic analysis showing co-segregation of the hypersensitive phenotype with the T-DNA insertion.

Genomic DNA from the mutant plants is isolated as described by Dellaporta et al, Plant Mol Biol Reporter 1: 19-21, 1983. A fragment of genomic DNA flanking the right border of the inserted T-DNA is rescued essentially according to Bouchez et al, Plant Mol Biol Reporter 14: 115-123, 1996, with minor modifications. Genomic DNA is digested with PstI, ethanol precipitated and resuspended in H₂O. DNA of vector pResc38 (Bouchez et al supra) is digested with PstI and dephosphorylated with shrimp alkaline phosphatase. The phosphatase is heat inactivated, the vector DNA is ethanol precipitated and resuspended in H₂O. 2.5 µg of PstI digested genomic DNA and 2.5 µg of PstI digested and dephosphorylated vector are mixed and ligated overnight at room temperature in a total volume 100 µl in the presence of 10 units of T4 DNA ligase. The DNA of the ligation mixture is precipitated with ethanol, resuspended in 50 µl H₂O, and digested with XbaI in a total volume of 100 µl. XbaI digested DNA is precipitated with ethanol and resuspended in H₂O. A second overnight ligation reaction in the presence of 10 units T4 DNA ligase is performed in a total volume of 200 µl at room temperature to achieve circularization of DNA fragments. The DNA of the ligation mixture is again precipitated with ethanol, rinsed two times with 70% ethanol, dried and dissolved in 5 µl H₂O. Two 2 µl aliquots are used for electroporation of electrocompetent E.coli XL1-Blue cells (Stratagene) according to the manufacturer's instructions. Transformants containing the inserted T-DNA and adjacent *Arabidopsis* genomic DNA sequences are selected on plates with 50mg/l kanamycin. Single bacterial colonies are analyzed by isolation of plasmid DNA using QIAprep Spin Plasmid Kit (Qiagen) and restriction digestion with PstI and XbaI. Plasmid pBN1 containing 3.7 kb of inserted T-DNA linked to 5 kb of *Arabidopsis* DNA is identified. Sequencing of the junction site is performed using a primer directed towards the flanking plant DNA and having the nucleotide sequence 5' -GGTTTCTACAGGACGTAACAT-3' (SEQ ID NO: 14) complementary to T-DNA 41 nucleotides from the right border. Digestion of this clone with SstI allows isolation of a 960 bp fragment which when labelled with ³²P can be conveniently used as a probe to screen wild type *Arabidopsis* genomic and cDNA libraries in order to identify the wild type gene affected in the *mkp1* mutant line.

Example 2: Cloning of the *AtMKP1* wild-type gene

The 960 bp *Sst*I fragment mentioned at the end of example 1 is labeled with ^{32}P by random oligonucleotide-primed synthesis (Feinberg et al, Anal Biochem 132: 6-13, 1983) for use as a probe in the following hybridization experiments.

Southern blot analysis of *Arabidopsis* wild type and *mkp1* DNA digested with *Eco*RV confirms that in the *mkp1* genomic DNA the sequence hybridizing to the probe is linked to T-DNA.

Northern blot analysis of *Arabidopsis* wild type RNA reveals the presence of a hybridizing transcript in RNA extracted from seven-day-old wild type seedlings. No such hybridizing fragment is detected in the corresponding RNA of *mkp1* seedlings.

A cDNA library (Elledge et al, 1991) and a genomic library (Stratagene) of wild type *Arabidopsis thaliana* ecotype Columbia is screened with the labelled *Sst*I fragment mentioned above. Screening of the bacteriophage λ libraries is performed according to the protocols described in chapter 6 of Ausubel et al, 1994, "Current protocols in molecular biology", John Wiley & Sons, Inc. Hybridization is performed as described by Church and Gilbert, Proc Natl Acad Sci USA 81: 1991-1995, 1984. Bacteriophage clones hybridizing to *Sst*I fragment are subjected to *in vivo* excision of plasmids according to Elledge et al, Proc Natl Acad Sci USA 88: 1731-1735, 1991, and Stratagene protocols. Inserts of the plasmids obtained are further analyzed by sequencing.

By partial sequencing and alignment of ten overlapping clones (pBN5.1 to pBN5.10) isolated from the genomic library a continuous genomic sequence of 6356 bp (see SEQ ID NO: 1) is decoded.

Ten cDNA clones representing the same gene, one of them a 3.0 kb full-length cDNA (SEQ ID NO: 2), are isolated from the cDNA library.

Example 3: Sequence Analysis and Alignments

The 3 kb full-length cDNA clone of SEQ ID NO: 2 encodes an ORF with the start codon being defined by basepairs 298-300 and the stop codon by basepairs 2650-2652. The ORF encodes a protein consisting of 784 amino acids (SEQ ID NO: 3) and a predicted molecular mass of 86.0 kD. Alignment with the genomic sequence of SEQ ID NO: 1 reveals three introns. T-DNA is inserted within the coding sequence of the *mkp1* mutant DNA before basepair position 502 according to the numbering of SEQ ID NO: 2. The sub-sequence

VHCCQGVSR (SEQ ID NO: 4) found in SEQ ID NO: 3 can be interpreted as corresponding to the mammalian sequence motif IHXAGXXRS (SEQ ID NO: 5) defining the family of protein tyrosine phosphatases, wherein the Ile at the first position can be replaced by Val and the Ser at the last position can be replaced by Thr (Van Vactor et al, Curr Opin Gen Dev 8: 112-126, 1998). Therefore it is concluded that the wild type ORF encodes a protein tyrosine phosphatase with invariant aspartic acid, cysteine, and arginine residues (Fauman et al, Trends Biochem Sci 21: 413-417, 1996) in positions 204, 235 and 241 according to SEQ ID NO: 3.

A data base search using the TFASTA program (Wisconsin Package Version 9.1, Genetics Computer Group (GCG), Madison, Wisc.) reveals that the encoded phosphatase has a significant similarity to dual specificity phosphatases. The closest homologue identified is *Xenopus laevis* MAP kinase phosphatase (MKP; Lewis et al, J Cell Sci 108: 2885-2896, 1995) showing 38.1% identity and 52.5% similarity in a 140 amino acid overlap region. The deduced AtMKP1 protein also has 36.0% identity and 52.5% similarity with a 140 amino acid overlap region encoded by the rat 3CH134/CL100 cDNA representing a rat MKP1. Essentially identical results are obtained when using the BLASTP 2.0.4 (Feb-24-1998) program of the NCBI BLAST family of programs which, allowing gapped alignment, compares an amino acid query sequence against a protein sequence database (Altschul et al, Nucleic Acids Res. 25: 3389-3402, 1997). No higher plant homologues are identified. The genomic position of the *AtMKP1* gene is determined by hybridization to filters containing genomic YAC clones publicly available from the Arabidopsis Biological Resource Center (Ohio, USA). The *AtMKP1* gene is found to map to chromosome 3 between markers ve022 and BGL1.

Example 4: **Complementation**

mkp1 mutant plants are transformed with DNA comprising the corresponding wild type genomic DNA including promoter and polyadenylation signal to find out whether the cloned wild type gene is able to complement the mutant *mkp1* phenotype.

mkp1 mutant plants harbor T-DNA containing the *NPTII* and *bar* marker genes under the control of nos and CaMV35S promoters, respectively. Therefore, different marker genes are used for the transformation construct. The vector used is a derivative of p1'barbi which is highly efficient in *Arabidopsis* transformation (Mengiste et al, Plant J 12: 945-948, 1997). In

p1'barbi the EcoRI fragment containing 1'promoter, *bar* gene coding region, and CaMV 35S polyadenylation signal is inverted in respect to the T-DNA borders by EcoRI digestion and re-ligation. In the resulting plasmid the 1'promoter (Velten et al, EMBO J 3: 2723-2730, 1984) is directed towards the right border of the T-DNA. This plasmid is digested with BamHI and NheI, and the *bar* gene and CaMV 35S polyadenylation signal are replaced by a synthetic polylinker with the sites for the restriction enzymes BamHI, HpaI, ClaI, StuI and NheI. The resulting plasmid is digested with BamHI and HpaI and ligated to a BamHI-PvuII fragment of pROB1 (Bilang et al, Gene 100: 247-250, 1991) containing the hygromycin-B-resistance gene *hph* linked to the CaMV 35S polyadenylation signal. The T-DNA of the resulting binary vector p1'hygi contains the hygromycin resistance selectable marker gene under the control of the 1'promoter and unique cloning sites for the restriction enzymes *ClaI*, *StuI* and *NheI* located between the marker gene and the T-DNA right border. p1'hygi is used to insert the reconstructed *AtMKP1* gene as follows. Plasmid pBN1 of example 1 is digested with *PstI* and *MunI* and dephosphorylated. The restriction fragment containing the 3'portion of the *AtMKP1* gene and pBluescript-SK(+) is purified from the agarose gel and ligated to the *PstI*-*MunI* restriction fragment of the wild type genomic clone pBN5.2 (example 2) including the 5' end of the coding sequence of the *AtMKP1* gene and 2.4 kb of upstream sequences. The reconstructed *AtMKP1* gene is excised by *PstI* and *NotI* and after filling the ends is inserted into the *StuI* site of p1'hygi. The construct is introduced by transformation into *Agrobacterium tumefaciens* strain C58CIRif^R containing the non-oncogenic Ti plasmid pGV3101 (Van Larebeke et al, Nature 252: 169-170, 1974). T-DNA containing the reconstructed *AtMKP1* gene is transferred to mutant plants by the method of *in planta Agrobacterium* mediated gene transfer (Bechtold et al, C R Acad Sci Paris, Life Sci 316: 1194-1199, 1993). Seeds of infiltrated plants are grown on hygromycin-containing medium to screen for transformants. The progeny of selfed hygromycin resistant plants is analyzed for the segregation of hygromycin resistance. The families in which a 3:1 segregation ratio is observed are used to isolate homozygous lines bearing the newly introduced T-DNA inserted at a single genetic locus. The obtained hygromycin resistant lines are analyzed by Northern blot analysis for the restoration of *AtMKP1* expression. In these lines the restoration of transcription of the *AtMKP1* gene can be observed as well as the restoration of the wild type level of MMS resistance and ABA mediated stress responses. Complementation is not observed in plants transformed with p1'hygi only.

Example 5: Cloning of homologous sequences from other plant species

Use of *AtMKP1* cDNA as a probe for Southern hybridization with genomic DNA from other plant species such as *Sinapis alba* (mustard), *Lycopersicum esculentum* (tomato) and *Zea mays* (maize) is successful in the case of *Sinapis alba* which belongs to the same family as *Arabidopsis* (Brassicaceae).

Homologous sequences from the other species can be identified in a PCR approach using degenerate primers 1-3 below, wherein I is inosine, derived from the regions conserved between VH-PTP13 of *Clamydomonas eugametos* and *AtMKP1* protein:

Primer 1 (forward): 5' -AAY AAY GGI ATH ACI CAY ATH YT-3' (SEQ ID NO: 11);

Primer 2 (reverse): 5' -YTG RCA IGC RAA ICC CAT RTT IGG-3' (SEQ ID NO: 12);

Primer 3 (reverse): 5' -IGT CCA CAT IAR RTA IGC DAT IAC (SEQ ID NO: 13);

A PCR reaction is performed in a total volume of 50 µl containing 1x reaction buffer (Qiagen), 200 µM of each dNTP, 1.25 units of Taq polymerase (Qiagen), and 100 pmol of each primer.

Reaction 1 is performed with primers 1 and 2 using genomic DNA from *Sinapis alba* (200 ng), *Lycopersicum esculentum* (400 ng), or *Zea mays* (600 ng) as the original template DNA. Amplification is carried out after an initial denaturation step of 3 min at 94°C, followed by 30 cycles of 30 sec at 94°C, 30 sec at 40°C, and 3 min at 72°C. The resulting amplification mixture is diluted 10³ fold.

Reaction 2 is performed using 2µl of the above dilution to provide the necessary template DNA. This time primers 1 and 3 are used under the same conditions as specified for reaction 1. The resulting amplification products are cloned into the T/A vector pCR2.1 (Invitrogen) and further analyzed by nucleotide sequencing.

Using this PCR approach it is possible to amplify sequences homologous to the *AtMKP1* gene from all the species mentioned above. Whereas the nucleotide sequence from *Sinapis alba* SaMKP1 ((SEQ ID NO: 15 encoding SEQ ID NO: 16) is 90.8% identical to the *AtMKP1* sequence, the nucleotide sequence from *Lycopersicum esculentum* LeMKP1 (SEQ ID NO: 17 encoding SEQ ID NO: 18) is 72.3% and the *Zea mays* sequence

ZmMKP1 (SEQ ID NO: 19 encoding SEQ ID NO: 20) 71.8% identical. The fragments hybridize to genomic DNA from corresponding species under the usual hybridization conditions for Southern blot analysis. The fragments can be used as probes to screen cDNA libraries for corresponding cDNA sequences.

The 243 bp *ZmMKP1* fragment amplifying from maize DNA is used as a probe to screen a maize cDNA library (Clontech) made in the Lambda ZAP[®]II Vector (Clontech) from "Blizzard" hybrid etiolated shoots, which were treated with the herbicide safener Benoxacor. The titer of the library is determined as 3×10^9 pfu/ml.

Library screening is conducted as described in the Clontech Lambda Library Protocol Handbook, with some slight modifications. Briefly, a single colony of XL-1 Blue is picked and incubated overnight at 37°C in LB medium, containing 10mM MgSO₄ and 0.2% maltose. 600 µl of stationary phase grown bacteria for each 150 mm plate is combined with 100 µl of phage library dilution in sterile 1× lambda dilution buffer (100mM NaCl; 10mM MgSO₄; 35mM Tris-HCl, pH7.5) to yield approximately 30,000 pfu per plate. This mixture is incubated at 37°C for 15 minutes. subsequently 7 ml of melted LB soft top agarose (at 48°C) is added to the cell suspension for each 150 mm plate, shortly mixed and then poured on two-day-old LB^{MgSO₄} agar plates, which have been pre-warmed to 37°C for four hours. The plates are then incubated at 37°C until plaques reach appropriate sizes (after about 8 to 9 hours). After chilling the plates at 4°C for one hour, phage particles are transferred to Hybond N nitrocellulose filters and the orientation of each filter to its plate is recorded with a waterproof pen. The filters are then treated by placing them on Whatman 3MM paper saturated with the appropriate solution. The treatments include denaturation solution (0.5M NaOH; 1.5M NaCl) for 2 minutes, followed by neutralization solution (0.5M Tris-HCl, pH 7.2; 1.5M NaCl; 1mM EDTA) for 3 minutes and 2× SSC for 3 minutes. DNA is subsequently crosslinked to the filters by UV.

Filters are then pre-hybridized, hybridized with the radioactively labeled *ZmMKP1* fragment and washed as described in Sambrook et al, Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press, chapters 9.47-9.57 and 11.45-11.49, 1989. An agar plug from the position of a positive plaque is then removed from the master plate and incubated overnight at 4°C in 1 ml of 1× lambda dilution buffer, containing 20 µl of chloroform. Each titer is determined and the phages are re-plated to obtain approximately 200 to 500 plaques on a 150 mm plate for a secondary screen as described above. Single plaques of interest

are collected from the agar plates and incubated over night at 4°C in 500 µl of 1× lambda dilution buffer and 20 µl of chloroform.

The pBluescript phagemid is excised from the λZAP™ vector as described by the *In Vivo* Excision Protocol using the ExAssist/SOLR System in the Stratagene Uni-ZAP™ XR Library Instruction Manual (1993). A 1/100 dilution is made of XL1-Blue MRF' and SOLR overnight cultures (at 30°C) and incubated at 37°C for 2-3 hours. XL1-Blue MRF' cells are then pelleted for 10 minutes at 1,500×g and re-suspended at an OD₆₀₀ = 1.0 in 10mM MgSO₄. 200 µl of these XL1-Blue cells are then combined with 250 µl of phage stock and 1 µl of ExAssist helper phage in a 50 ml conical tube and incubated at 37°C for 15 minutes. 3 ml of LB broth is added and incubated at 37°C for 5 hours, after which the cells are pelleted for 15 minutes at 2,000×g and the supernatant transferred to a new tube. The tube is then heated at 70°C for 15 minutes and centrifuged again for 15 minutes at 4,000×g. The supernatant, containing the excised phagemid pBluescript packaged as filamentous phage particles, is decanted into a new tube. 10 and 100 µl of this phage stock are then added to two tubes with 200 µl of SOLR cells that have been allowed to grow to OD₆₀₀ = 0.5-1.0 before being removed from the incubator and further incubated at room temperature. The tubes are incubated at 37°C for 15 minutes, followed by plating 10-50 µl from each tube on LB^{amp} (50 µg/ml) and over night incubation at 37°C.

The positive clones are checked for insert size by *EcoRI/XhoI* double digestion and end-sequencing with T3 and T7 promoter primers (Promega).

Screening of 360,000 pfu of the library results in three identical clones of 2.2 kb containing the 3' poly(A) tail, but lacking part of the 5' end, including the translation initiation site. The gene corresponding to the identified partial cDNA clone is named *ZmMKP2*, as it is not identical with the *ZmMKP1* fragment used as the probe (92.3% identity on the nucleotide level over the 196 bp fragment flanked by the primers 1 and 3). An additional 213 nucleotides are amplified and cloned by 5' RACE (rapid amplification of cDNA ends) carried out following the instructions of the 5'/3' RACE Kit (Boehringer Mannheim) resulting in a longer cDNA sequence of 2,452 bp but still not complete, judged by the predicted mRNA length from the RNA gel blot analysis and the absence of a possible translation initiation site.

The sequence information gained from the *ZmMKP2* cDNA including the additional 213 nucleotides obtained by 5' RACE (SEQ ID NO: 21 encoding SEQ ID NO: 22) is used to

design two additional backward oriented degenerate primers wherein I is inosine to 3' regions conserved between the deduced peptide sequences of *ZmMKP2* and *AtMKP1*:

Primer 4 (reverse): 5'-GCI GCY TTI GCR TCY TTY TCC-3' (SEQ ID NO: 25);

Primer 5 (reverse): 5'-YTC ICK IGC IGG IAR RTG IGT YTC-3' (SEQ ID NO: 26)

These primers are used to PCR amplify a larger fragment of a MAP kinase phosphatase gene from tomato. The amplified and cloned 522 bp long fragment is not identical to *LeMKP1*. Therefore, its corresponding gene is named *LeMKP2* (SEQ ID NO: 23 encoding SEQ ID NO: 24; 75% identity on the nucleotide level over the stretch of 196 bp of *ZmMKP1* flanked by primers 1 and 3). The origins of all identified MAP kinase phosphatase homologous gene sequences are confirmed by Southern blot analysis.

The following Table shows an alignment of a continuous stretch of 312 amino acids of *AtMKP1* with the related amino acid sequence of *ZmMKP2*.

AtMKP1	139	KREKIAFFDKECSKVADHTYVGDAVAKDKSILKNNGITHILNCVGFICP	188
		:::. : : : . : . : : :	
ZmMKP2	24	RKDQIAFFDKECSKVADHVYLGDAVAKNRDILLRNGITHVLNCVGFVCP	73
AtMKP1	189	EYFKSDFCYRSLWLQDSPSEDITSILYDVFDYFEDVREQSGRIFVHCCQG	238
		. . : : :	
ZmMKP2	74	EYFKSDLVYRTLWLQDSPTEDITSILYDVFDYFEDVREQGGRVLVHCCQG	123
AtMKP1	239	VSRSTSLVIAYLMMREGQSFDDAFQYVKSARGIADPNMGFACQLLQCQKR	288
		: : : : . :	
ZmMKP2	124	VSRSTSLVIAYLMMREGQSFDDAFQFVKAARGIANPNMGFACQLLQCQKR	173
AtMKP1	289	VHAFPLSPTSLLRMVYKMSPHSPYDPLHLVPKLLNDPCPGSLDSRGAFIIQ	338
		. : . : . :	
ZmMKP2	174	VHAIPLSFNSVLRMYRMAPHSQYAPLHLVPKMLNDPSPATLDSRGAFIVH	223
AtMKP1	339	LPSAIYIWVGRQCETIMEKDAKAAVCQIARYEKVEAPIMVREGDEPVYY	388
		. .: : . : : : :	
ZmMKP2	224	VLSSLYVWGMKCDPVMEKDAKAAAFQVRYEKVQGHIVVREGLEPQEF	273
AtMKP1	389	WDAFASILPMIGG.....SVIKVQPGDRKVDAYNLDFEIFQKAIE	428
		.	
ZmMKP2	274	WDAFSSMPNDSNTKISKDQIDSASKSDPGSRKNESYDADFELVYKAIT	323
AtMKP1	429	GGFVPTLASSNNEHETHLPARE	450
		..	
ZmMKP2	324	GGVVPFSTSGAGDETHLPARE	345

What is claimed is:

1. A DNA comprising an open reading frame encoding a protein characterized by a component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO: 3
2. The DNA according to claim 1 comprising an open reading frame encoding a plant MAP kinase phosphatase.
3. The DNA according to claim 1 wherein the open reading frame encodes an amino acid sequence selected from the group of amino acid sequences described in SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.
4. The DNA according to claim 1, wherein the open reading frame encodes a protein characterized by the amino acid sequence of SEQ ID NO: 3
5. The DNA according to claim 1 characterized by the nucleotide sequence of SEQ ID NO: 1
6. The DNA according to claim 1 wherein the open reading frame encodes a protein contributing to repair of DNA damage in a plant cell.
7. The DNA according to claim 1 wherein the open reading frame encodes a protein conferring hypersensitivity to treatment with methyl methanesulfonate (MMS).
8. The DNA according to claim 7 wherein the open reading frame encodes a protein conferring hypersensitivity to treatment with UV light or X-rays.
9. The DNA according to claim 7 wherein the open reading frame encodes a protein interfering with abscisic acid signal transduction.
10. The protein encoded by the open reading frame of any one of claims 1 to 9.
11. A method of producing DNA according to claim 1, comprising
 - screening a DNA library for clones which are capable of hybridizing to a fragment of the DNA defined by SEQ ID NO: 1, wherein said fragment has a length of at least 15 nucleotides;
 - sequencing hybridizing clones;
 - purifying vector DNA of clones comprising an open reading frame encoding a protein with more than 40% sequence identity to SEQ ID NO: 3
 - optionally further processing the purified DNA.
12. A polymerase chain reaction wherein at least one oligonucleotide used comprises a sequence of nucleotides which represents 15 or more basepairs of SEQ ID NO: 1.

SEQUENCE LISTING

<110> Novartis AG

<120> Organic compounds

<130> Silencing Gene

<140>

<141>

<160> 26

<170> PatentIn Ver. 2.1

<210> 1

<211> 6356

<212> DNA

<213> Arabidopsis thaliana

<400> 1

```

gttcaagggtg gtgttatttn cagtattaga aaagaggcctt ctagagagag tttctaaatt 60
atattttgaa cacggccgga ttggttgtct tatgattaag tgaatgcctt agatctgggtg 120
acgattggta tatgagatat atagtanaat gatagaacat cagaaatact ataagtcacc 180
atatttttaa aaaataaatg ctatagaatt tttgttttgg taattgttat aactctacaa 240
agttatgtgt tatagagttt tactagtttc atcgttatca tgtgtatagt ataacncaac 300
aaaagaaatt ttaatatctt gaaaacataa aaatttaata aaatgatgtg agtataagaa 360
aaaaagaaag aaagaaacaa cgtaaaaaat aaaaatcatt catacatata acaattttca 420
aaagatcaat gttaacttta attagtcttt tttcttatgt ttcattgcaa tgcaatagta 480
ttacttttct ttaatctaag attatgtgtt gcttttaagc aagaactatt caaagagtca 540
aagacatgca tgtaaaacttt agagaacggg atctttcatc aatcttattt ttttcccttt 600
tttttttaca acgaacgata ggtagtaagc atgaatctgg ttcaaagttt ggatcagtgt 660
gggcataatc ctggttcgct aaaaagaatc aatataatat atgttgagcg taagagaaga 720
aagctcatag ttctgtgtag aaaaaacgtg gagattggag agaaattact aagagagcga 780
agagaagagt aaacctagaa gatacaaaag acgctatcaa aagattgttt tatgtgttta 840
tgaacatttc acttaggcca gtataaatat taaatgtcat tttgagaaag aaaaaaaaaa 900
ggtgtcattt tgaggatata aatattttct caacaagatg ttttagactt tcaacataac 960
gatcatttaa aactacataa tttatctctt cgttaaaaaa ttaacattaa caaaaagtta 1020
aattcgacaa gaagagttaa ttcaataagt atactgtatg tgtttatttt attcacatga 1080
aaagtgtcaa acaacacttg ggatacatat ttcattgcta aaaatcgat aacaacattt 1140
aaggtaatca gtaaatattg tgcctaaact aaaatatgtg accattttga ccaatgagca 1200
ttttttggaa ataaaactac tgcatacgac ctgggtcaaac acatgaactg ccactgcttt 1260
taaaactacc tttttaaaact tttgacgtcc atttttattt actaattatt atgttaatca 1320
aaccactaac attatgattc gaaattttcg gtgatgattg ttagaagtga tgtgaatgat 1380
gcggtataaa acaaatgttg atatatattc actcatatat atcaaaatca aaatagctta 1440
tcgttccaaa acatgattga taaaatgtaa ttactattca aaacttaaag ccagtagtta 1500
aaataaaaca aaataaatag tatatgtttc atattaatgc ccaaccaata ttttgttttg 1560
atgaggaaaa gtcttttttt ttttttcttt ttttttctca tagttaaaag acataaaaaa 1620
aattaatatt acaacaaga caaaaaaaa gaataaataa aacaacaatc attactgtca 1680
aatcaattat cacagggaaa agttattaag aaatgtcaac caatgagtat cattatcatc 1740
atgctttttt agacccttct ttttaattca tcaagattta gtcttgttta taaatttgaa 1800
gtaaaaattc aaatgaatca attctacaat tttttcccta cattattgta acgataaaat 1860
ttaagatcaa caattacttc gtttaatttt tttctgataa aaatatttga tatctttctg 1920
attatgatat attagcattg tttttgtatt gtatgtatgg taattaattt tagttcaaaa 1980

```


gaataataaa	tgggttgatt	agcatgaatt	taataaaaaa	ataagactga	ataaacatag	2040
gtaataaaact	ttgttttcttt	tggtaaagt	aaaattagaa	aaaatcataa	tcccaaggta	2100
attactataa	ttacattcaa	tgtcagaatt	aaacgtagt	aataaaaaacc	gtaaaacatg	2160
agaaaaacaa	gaatttatta	tctttgacaa	gcaataaatg	aatgctgac	aaaaaattgg	2220
tttcaaagtt	tcaacgcgtt	tcttataata	agaattcaat	ttcgtgcagc	taatcaggag	2280
ataattatca	taattaaatt	aatcgttact	actttataat	actcccaaaa	taatcgaaaa	2340
cgaatttatt	ttattgtaat	ttgtttaata	ataaagaatt	actgtttcct	cccacttccc	2400
atctctcttt	tctttttgtg	ttcttctctt	tctccgcttt	gtttcccaaa	tctctctctt	2460
ctctctctct	gaagaaaaat	aaataaaaaga	tctaactttg	acggctctct	taatcttact	2520
cactccgtaa	gttccaaatc	tctctccttt	actctcatat	ctatatcgtc	cgaacaaaac	2580
ccaggagaat	tgcttcaccc	cctttttggg	tttttaataca	ttttctcaga	ttctcagttt	2640
ctgtttccgt	cttctagatt	ctgggttcag	tttctgtttt	gctcttattg	aattttctta	2700
ttcattttgt	gtttcggagt	tattcatggg	agctgaattt	gttaattctg	atgttggttt	2760
gcgttttctt	cctttctagt	ttggctatgt	cgtctttgat	ctgatgctgg	gttattctct	2820
ttccctctgt	tttggtttct	tttaggggtt	taagtccgaa	tagactgatg	ggagcttgat	2880
ggttattggt	agatcagatg	tggatttaaa	gccttcgctg	aactaacaag	tctatggaag	2940
aagcaaagac	ccttggttta	cactgtatgt	tgtgaggaat	ttgtctgatt	ttgggtgata	3000
aaggtgaagt	ccttgagttt	gtaattttga	gataagattg	gatgggtgga	agagaggatg	3060
cgatggggaa	tgatgaagct	cctcctgggt	ctaagaaaat	gttttgccgg	tctgcctctt	3120
ggtctgcttc	acggactgca	tcacaagttc	ctgaggggtga	tgagcaaaagc	ctgaacattc	3180
cgtgtgctat	tagttctggg	ccgagtcgaa	gatgtccagc	tgctcctttg	acacctcggt	3240
cacatcataa	cagcaaggct	agagcttggt	tgccaccatt	gcagcctctt	gccattttcta	3300
ggaggagctt	agacgagtgg	cctaaggcgg	gttcggatga	tgtcggtgag	tggcctcatc	3360
caccaacacc	tagcgggaac	aaaaccgggg	agagattgaa	gctcgattta	tcatacaacgc	3420
agcagcgggt	aacagataag	agctctgggt	tagctaagag	ggagaagatt	gctttctttg	3480
ataaagaatg	ttcaaagggt	gctgatcata	tatatgtggg	tggaagtgct	gtggcgaaag	3540
acaagagcat	actgaaaaac	aatggaatca	cgcatactct	gaactgcgtt	ggttttatct	3600
gtccggaata	tttcaagttc	gattttttgt	acagatcctt	gtggttacag	gatagtccgt	3660
cagaggatat	agctagtatt	ctgtacgatg	tgtttgacta	ccttgaagat	gtgagggagc	3720
aaagtggaag	gatctttgtt	cattgttggtc	aaggggtttc	acgatctacc	tcgttggttaa	3780
tagcatatct	gatgtggaga	gaagggcaaa	gttttgatga	tgcatctcag	tatgtgaagt	3840
ctgctagagg	tattgctgat	cctaacatgg	gctttgcttg	ccaattgtta	caatgccaaa	3900
agaggggttca	tgcgttcccc	cctagcccta	cctccttact	tagaatgtac	aaaatgtctc	3960
cacactctcc	ttatgacctt	ttgcatcttg	ttccaaaact	gttgaatgat	ccatgcccca	4020
gttctctgga	ttcaagaggt	gcatttatca	ttcagttacc	ttctgcaatt	tacatttggg	4080
ttggtaggca	gtgtgaaacc	atcatggaga	aagatgcaaa	agctgctggt	tgtcagattg	4140
ctcgatatga	gaaagtcgaa	gcacctatta	tgggtggtcag	agaaggtgat	gagcctgttt	4200
attactggga	tgcatttgca	agcattttgc	ctatgatgtg	gggctcggtta	attaaagtct	4260
aaccagggtga	caggaaaggtc	gacgcataata	atctggattt	tgagattttt	cagaaagcca	4320
tagagggag	ttttgtgcca	actttagcat	catccaacaa	cgaacatgag	actcatcttc	4380
ctgcaaggga	aaacagttgg	agctcactga	aatgtaagtt	tgcatcaagg	tttgacaaag	4440
gttttccgga	tgtctccaaa	acgccactat	ctaggggtcta	ttcagattcg	atgatgatcg	4500
tgcatctcat	aggctcacct	tcttcaacaa	ccttctcctc	atccactgct	tcgcctcctt	4560
ttctctctct	cgattctgta	tggttcaacaa	attcaggcaa	tagcttaaag	agtttctctc	4620
aatcctctgg	acgttcgtcc	ttgagacctt	ctattccacc	atcgctgaca	ttgcctaaat	4680
tttccagcct	atccctcctc	ccttcccaaa	ccttctcctaa	agaatctcgt	ggtgtcaata	4740
cctttcttca	accgtcacca	aatagaaaagg	cctcaccttc	tcttgctgag	cgtagaggca	4800
gcctgaaagg	atctctgaag	ttgccagggt	tgggtgatcc	caacagaggc	acacctgctt	4860
ttactttaca	tccggatgat	agtaatgaca	tagtcttcaa	tctggagggt	attagaaacg	4920
gcgatctata	tccaccaagt	gattgcaaa	ggacaagtgt	agattcagat	ttgccagaga	4980
aggaaattat	atccttaatc	agttgcagta	aatctgacag	acacaaatcg	ggagggtgata	5040
ctgatagctc	tggccagcct	ttagcatgtc	gttggccaag	tatggagatg	attacaaaac	5100
tgagcagagc	ttacttagat	tcagaatctg	ttatagcaat	cccgttgcca	agtgatgctg	5160
taggagaaac	gggttagtagg	aatttgtaca	tttggatcgg	aaagtcattc	tctttggata	5220
acaactgttc	tctagtagat	agcaacaaag	cggcagacac	tgtggagaat	gttgattggg	5280

tacaaattgg	tgaatccatt	ttgtgtcaga	tggacttgcc	aaaagatacc	cctataaagg	5340
taataatagc	ctaaactttg	gaggctctga	tactttttac	taattgtaaa	gtctgcgtgc	5400
tcatctttgt	catgtcttat	ccaaccaaac	tatatttcga	agatgaaaat	tacaatctca	5460
gcactttcat	tactgactac	tgaggacggt	taggtagaat	ccttatgatt	tcagcagttg	5520
tatgtattgg	tttattctct	agtggtttgc	atggttccaa	cttggtatga	tccttttgtt	5580
gtttgtaact	gataagttgc	ttttctttct	tgtaaacaga	tagttaggga	atctgaggat	5640
cagacagagt	tgctagcact	gctgagcgcg	ctataacacc	cacccgcaag	ctctacacat	5700
ttactctgtt	tttttttcac	agattccttc	aaccgcaaca	cttttccatt	ttcagacaga	5760
gtattcattc	agctcaggtg	agaattctct	gaaagcagtc	tgtaacactt	catcttcaca	5820
gttgcacccg	aatacaatcg	ttagtctctg	attatgttta	attgctatct	gatcatgaat	5880
ttgagttaga	ggatgggttg	aacaaaaaaa	cttagaagct	cgaatgaccg	gtttttacca	5940
aattctcata	gaccatattt	gattcttttg	atttacttct	ggtgcaggac	tctctgtgct	6000
tatggaagtt	gatgttgggg	gaaacaactc	tctgttacag	tggggaaaaa	acttcttctt	6060
cttctttcta	tcacatgaaa	atcctcaagg	gccattatta	gtatgatcag	attataaaat	6120
tgtaagggtta	ggggctttat	gaggattttg	atggacttgt	tacaatgttt	acataacac	6180
tcagcagcac	aatagatttt	tgttaaactt	acatgttatt	caagtaaaag	tactatgtag	6240
atgttgaagt	ctaattgaag	aattagttaa	tgatagtctt	aaacacttga	ttcacttgtc	6300
atccaatttt	ggttttgcgc	atagtttctc	ttcttttatt	tcctctctaa	aacacg	6356

<210> 2

<211> 3059

<212> DNA

<213> *Arabidopsis thaliana*

<400> 2

cttctttctc	gctttgtttc	cccaatctct	ctcttctctc	tctctgaaga	aaaataaata	60
aaagatctaa	ctttgacggc	tctcttaatc	ttactcactc	cgggttttaag	tcggaataga	120
ctgatgggag	cttgatgggt	attgttagat	cagatgtgga	tttaaagcct	tcgctgaact	180
aacaagtcta	tggaagaagc	aaagaccctt	gttttacact	gtatgttgtg	aggaatttgt	240
ctgatttttg	gtgataaagg	tgaagtcttt	gagtttgtaa	ttttgagata	agattggatg	300
gtgggaagag	aggatgcgat	ggggaatgat	gaagctcctc	ctggttctaa	gaaaatgttt	360
tggcggtctg	cctcttggtc	tgcttcacgg	actgcatcac	aagttcctga	gggtgatgag	420
caaagcctga	acattccgtg	tgctattagt	tctgggccga	gtcgaagatg	tcagctgct	480
cctttgacac	ctcggttcaca	tcataacagc	aaggctagag	cttgtttgcc	accattgcag	540
cctcttgcca	tttctaggag	gagcttagac	gagtggccta	aggcgggttc	ggatgatgtc	600
ggtgagtggc	ctcatccacc	aacacctagc	gggaacaaaa	ccggggagag	attgaagctc	660
gatttatcat	caacgcagca	gcgggtaaca	gataagagct	ctggtctagc	taagagggag	720
aagattgctt	tctttgataa	agaatgttca	aaggttgctg	atcatatata	tgtgggtgga	780
gatgctgtgg	cgaaagacaa	gagcatactg	aaaaacaatg	gaatcacgca	tatcttgaac	840
tgcgttggtt	ttatctgtcc	ggaatatattc	aagtctgatt	tttgttacag	atccttgtgg	900
ttacaggata	gtccgtcaga	ggatataact	agtattctgt	acgatgtgtt	tgactacttt	960
gaagatgtga	gggagcaaag	tggaaggatc	tttgttcatt	gttgtcaagg	ggtttcacga	1020
tctacctcgt	tggtaatagc	atatctgatg	tggaagagaag	ggcaaagtgt	tgatgatgca	1080
tttcagtatg	tgaagtctgc	tagagggtatt	gctgaccta	acatgggctt	tgcttgccaa	1140
ttgttacaat	gccaaaagag	ggttcatgcg	ttcccgctta	gccctacctc	cttacttaga	1200
atgtacaaaa	tgtctccaca	ctctccttat	gaccttttgc	atcttgttcc	aaaactgttg	1260
aatgatccat	gcccggttcc	tctggattca	agaggtgcat	ttatcattca	gttaccttct	1320
gcaatttaca	tttgggttgg	taggcagtgt	gaaacatca	tggaagaaag	tgcaaaagct	1380
gctgtttgtc	agattgctcg	atatgagaaa	gtcgaagcac	ctattatggt	ggtcagagaa	1440
ggtgatgagc	ctgtttatta	ctgggatgca	tttgcaagca	ttttgcctat	gattgggggc	1500
tcggttaatta	aagttcaacc	aggtgacagg	aaggctgacg	catataatct	ggatttttag	1560
atttttcaga	aagccataga	gggagggttt	gtgccaaact	tagcatcatc	caacaacgaa	1620
catgagactc	atcttcctgc	aagggaaaaac	agttggagct	cactgaaatg	taagtttgca	1680
tcaagggttg	acaaagggtt	tccgtatgtc	tccaaaacgc	cactatctag	ggtctattca	1740
gattcgatga	tgatcgtgca	ttcatcaggc	tcaccttctc	caacaacttc	ttcatcatcc	1800

- 4 -

```

actgcgtcgc ctccttttct ctctcccgat tctgtatgtt caacaaattc aggcaatagc 1860
ttaaagagtt tctctcaatc ctctggacgt tcgtccctga gaccttctat tccaccatcg 1920
ctgacattgc ctaaattttc cagcctatcc ctctccctt cccaaacttc tcctaaagaa 1980
tctcgtggtg tcaatacttt tcttcaaccg tcaccaaata gaaaggcttc accttctctt 2040
gctgagcgta gaggcagcct gaaaggatct ctgaagttgc caggtttggc tgattccaac 2100
agaggcacac ctgctttttac ttacatccg gatgatagta atgacatagt cttcaatctg 2160
gaggggtatta gaaacggcga tctatatcca ccaagtgtt gcaaagggac aagtgtagat 2220
tcagatttgc cagagaagga aattatatcc ttaatcagtt gcagtaaatac tgacagacac 2280
aaatcgggag gtgatactga tagctctggc cagcctttag catgtcgttg gccaaagtatg 2340
gagatgatta caaaactgag cagagcttac ttagattcag aatctgttat agcaatcccg 2400
ttgccaaagt atgctgtagg agaaacgggt agtaggaatt tgtacatttg gatcggaag 2460
tcattctctt tggataacaa ctgttctcta gtatagatgca acaaagcggc agacactgtg 2520
gagaatgttg attgggtaca aattggtgaa tccattttgt gtcagatgga cttgccaaaa 2580
gataccccta taaagatagt tagggaatct gaggatcaga cagagttgct agcactgctg 2640
agcgcgctat aacacccacc cgcaagctct acacatttac tctgtttttt ttccacagat 2700
tccttcaacc gcaacacttt tccattttca gacagagtat tcattcagct caggactctc 2760
tgtgcttatg gaagttgatg ttgggggaaa caactctctt gtacagtggg gaaaaaactt 2820
cttctcttct tttctatcac atgaaaatcc tcaagggcca ttattagtat gatcagatta 2880
taaaattgta aggttagggg ctttatgagg attttgatgg acttgttaca atgtttacat 2940
atacactcag cagcacaata gatttttgtt aaacttacat gttattcaag taaaagtact 3000
atgtagatgt tgaagtctaa ttgaagaatt agttaatgat aaaaaaaaaa aaaaaaaaaa 3059

```

<210> 3

<211> 784

<212> PRT

<213> Arabidopsis thaliana

<400> 3

```

Met Val Gly Arg Glu Asp Ala Met Gly Asn Asp Glu Ala Pro Pro Gly
  1                      5                      10                      15

```

```

Ser Lys Lys Met Phe Trp Arg Ser Ala Ser Trp Ser Ala Ser Arg Thr
          20                      25                      30

```

```

Ala Ser Gln Val Pro Glu Gly Asp Glu Gln Ser Leu Asn Ile Pro Cys
          35                      40                      45

```

```

Ala Ile Ser Ser Gly Pro Ser Arg Arg Cys Pro Ala Ala Pro Leu Thr
          50                      55                      60

```

```

Pro Arg Ser His His Asn Ser Lys Ala Arg Ala Cys Leu Pro Pro Leu
          65                      70                      75                      80

```

```

Gln Pro Leu Ala Ile Ser Arg Arg Ser Leu Asp Glu Trp Pro Lys Ala
          85                      90                      95

```

```

Gly Ser Asp Asp Val Gly Glu Trp Pro His Pro Pro Thr Pro Ser Gly
          100                      105                      110

```

```

Asn Lys Thr Gly Glu Arg Leu Lys Leu Asp Leu Ser Ser Thr Gln Gln
          115                      120                      125

```

```

Arg Val Thr Asp Lys Ser Ser Gly Leu Ala Lys Arg Glu Lys Ile Ala
          130                      135                      140

```

- 5 -

Phe Phe Asp Lys Glu Cys Ser Lys Val Ala Asp His Ile Tyr Val Gly
 145 150 155 160
 Gly Asp Ala Val Ala Lys Asp Lys Ser Ile Leu Lys Asn Asn Gly Ile
 165 170 175
 Thr His Ile Leu Asn Cys Val Gly Phe Ile Cys Pro Glu Tyr Phe Lys
 180 185 190
 Ser Asp Phe Cys Tyr Arg Ser Leu Trp Leu Gln Asp Ser Pro Ser Glu
 195 200 205
 Asp Ile Thr Ser Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val
 210 215 220
 Arg Glu Gln Ser Gly Arg Ile Phe Val His Cys Cys Gln Gly Val Ser
 225 230 235 240
 Arg Ser Thr Ser Leu Val Ile Ala Tyr Leu Met Trp Arg Glu Gly Gln
 245 250 255
 Ser Phe Asp Asp Ala Phe Gln Tyr Val Lys Ser Ala Arg Gly Ile Ala
 260 265 270
 Asp Pro Asn Met Gly Phe Ala Cys Gln Leu Leu Gln Cys Gln Lys Arg
 275 280 285
 Val His Ala Phe Pro Leu Ser Pro Thr Ser Leu Leu Arg Met Tyr Lys
 290 295 300
 Met Ser Pro His Ser Pro Tyr Asp Pro Leu His Leu Val Pro Lys Leu
 305 310 315 320
 Leu Asn Asp Pro Cys Pro Gly Ser Leu Asp Ser Arg Gly Ala Phe Ile
 325 330 335
 Ile Gln Leu Pro Ser Ala Ile Tyr Ile Trp Val Gly Arg Gln Cys Glu
 340 345 350
 Thr Ile Met Glu Lys Asp Ala Lys Ala Ala Val Cys Gln Ile Ala Arg
 355 360 365
 Tyr Glu Lys Val Glu Ala Pro Ile Met Val Val Arg Glu Gly Asp Glu
 370 375 380
 Pro Val Tyr Tyr Trp Asp Ala Phe Ala Ser Ile Leu Pro Met Ile Gly
 385 390 395 400
 Gly Ser Val Ile Lys Val Gln Pro Gly Asp Arg Lys Val Asp Ala Tyr
 405 410 415
 Asn Leu Asp Phe Glu Ile Phe Gln Lys Ala Ile Glu Gly Gly Phe Val
 420 425 430
 Pro Thr Leu Ala Ser Ser Asn Asn Glu His Glu Thr His Leu Pro Ala

- 6 -

435		440		445
Arg Glu Asn Ser Trp Ser Ser Leu Lys Cys Lys Phe Ala Ser Arg Phe				
450		455		460
Asp Lys Gly Phe Arg Tyr Val Ser Lys Thr Pro Leu Ser Arg Val Tyr				
465		470		475
Ser Asp Ser Met Met Ile Val His Ser Ser Gly Ser Pro Ser Ser Thr				
		485		490
Thr Ser Ser Ser Ser Thr Ala Ser Pro Pro Phe Leu Ser Pro Asp Ser				
		500		505
Val Cys Ser Thr Asn Ser Gly Asn Ser Leu Lys Ser Phe Ser Gln Ser				
		515		520
Ser Gly Arg Ser Ser Leu Arg Pro Ser Ile Pro Pro Ser Leu Thr Leu				
		530		535
Pro Lys Phe Ser Ser Leu Ser Leu Leu Pro Ser Gln Thr Ser Pro Lys				
		545		550
Glu Ser Arg Gly Val Asn Thr Phe Leu Gln Pro Ser Pro Asn Arg Lys				
		565		570
Ala Ser Pro Ser Leu Ala Glu Arg Arg Gly Ser Leu Lys Gly Ser Leu				
		580		585
Lys Leu Pro Gly Leu Ala Asp Ser Asn Arg Gly Thr Pro Ala Phe Thr				
		595		600
Leu His Pro Asp Asp Ser Asn Asp Ile Val Phe Asn Leu Glu Gly Ile				
		610		615
Arg Asn Gly Asp Leu Tyr Pro Pro Ser Asp Cys Lys Gly Thr Ser Val				
		625		630
Asp Ser Asp Leu Pro Glu Lys Glu Ile Ile Ser Leu Ile Ser Cys Ser				
		645		650
Lys Ser Asp Arg His Lys Ser Gly Gly Asp Thr Asp Ser Ser Gly Gln				
		660		665
Pro Leu Ala Cys Arg Trp Pro Ser Met Glu Met Ile Thr Lys Leu Ser				
		675		680
Arg Ala Tyr Leu Asp Ser Glu Ser Val Ile Ala Ile Pro Leu Pro Ser				
		690		695
Asp Ala Val Gly Glu Thr Gly Ser Arg Asn Leu Tyr Ile Trp Ile Gly				
		705		710
Lys Ser Phe Ser Leu Asp Asn Asn Cys Ser Leu Val Asp Ser Asn Lys				
		725		730
				735

- 7 -

Ala Ala Asp Thr Val Glu Asn Val Asp Trp Val Gln Ile Gly Glu Ser
 740 745 750

Ile Leu Cys Gln Met Asp Leu Pro Lys Asp Thr Pro Ile Lys Ile Val
 755 760 765

Arg Glu Ser Glu Asp Gln Thr Glu Leu Leu Ala Leu Leu Ser Ala Leu
 770 775 780

<210> 4

<211> 10

<212> PRT

<213> Arabidopsis thaliana

<400> 4

Val His Cys Cys Gln Gly Val Ser Arg Ser
 1 5 10

<210> 5

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: mammalian
 sequence motif defining the family of PTPs

<400> 5

Ile His Cys Xaa Ala Gly Xaa Xaa Arg Ser
 1 5 10

<210> 6

<211> 14

<212> PRT

<213> Arabidopsis thaliana

<400> 6

Thr Ser Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val
 1 5 10

<210> 7

<211> 12

<212> PRT

<213> Arabidopsis thaliana

<400> 7

Phe Val His Cys Cys Gln Gly Val Ser Arg Ser Thr

- 8 -

1	5	10
---	---	----

<210> 8
<211> 4
<212> PRT
<213> Arabidopsis thaliana

<400> 8
Phe Val His Cys
1

<210> 9
<211> 6
<212> PRT
<213> Arabidopsis thaliana

<400> 9
Gln Gly Val Ser Arg Ser
1 5

<210> 10
<211> 5
<212> PRT
<213> Arabidopsis thaliana

<400> 10
Tyr Phe Lys Ser Asp
1 5

<210> 11
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: degenerate
primer

<400> 11
aayaayggna thacncayat hyt

<210> 12
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: degenerate
primer

<400> 12

23

grcanger aannccatrt tngg 24

<10> 13
 <11> 24
 <12> DNA
 <13> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: degenerate primer

<400> 13
 ngtcacatn arrtangcda tnac 24

<210> 14
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 14
 ggtttctaca ggacgtaaca t 21

<210> 15
 <211> 196
 <212> DNA
 <213> Sinapis alba

<400> 15
 caactgcgtt ggtttcatct gtccctgaata tttcaagtct gatttttggtt accgggtcggtt 60
 gtgggttacgt gatagtccat cagaggatat aactagcatt ctctacgatg tctttgacta 120
 ctttgaagac gttagggagc aaagtgggag gatctttggt cactgttggtc aaggcggttc 180
 acggtctacc tccttg 196

<210> 16
 <211> 65
 <212> PRT
 <213> Sinapis alba

<400> 16
 Asn Cys Val Gly Phe Ile Cys Pro Glu Tyr Phe Lys Ser Asp Phe Cys
 1 5 10 15
 Tyr Arg Ser Leu Trp Leu Arg Asp Ser Pro Ser Glu Asp Ile Thr Ser
 20 25 30
 Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val Arg Glu Gln Ser
 35 40 45
 Gly Arg Ile Phe Val His Cys Cys Gln Gly Val Ser Arg Ser Thr Ser
 50 55 60
 Leu

- 10 -

65

<210> 17
 <211> 196
 <212> DNA
 <213> Lycopersicon esculentum

<400> 17
 aaactgtgta gggtttagtt gtcctgaata ctttaaggat gaccttgtat acaagacact 60
 ttggctgcag gatagcccca ctgaggacat cacaagtatt ctttatgatg tctttgatta 120
 ctttgaagat gttcatgaac aagggtgggag tgtctttgta cactgcttcc aggggggtgct 180
 ccgatcagcc tccttg 196

<210> 18
 <211> 65
 <212> PRT
 <213> Lycopersicon esculentum

<400> 18
 Asn Cys Val Gly Phe Ser Cys Pro Glu Tyr Phe Lys Asp Asp Leu Val
 1 5 10 15
 Tyr Lys Thr Leu Trp Leu Gln Asp Ser Pro Thr Glu Asp Ile Thr Ser
 20 25 30
 Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val His Glu Gln Gly
 35 40 45
 Gly Ser Val Phe Val His Cys Phe Gln Gly Val Ser Arg Ser Ala Ser
 50 55 60

Leu
 65

<210> 19
 <211> 196
 <212> DNA
 <213> Zea mays

<400> 19
 caactgcatg ggcttcgtct gcccgaagta cttcaagtct gaccttgtct accgcaccct 60
 ctggctgcag gacagcccca ccgaggacac caccagcatc ctttacgacg tgtttgatta 120
 ctttgaggac gtcagggagc aggggtggccg cgtgtttgtg cattgctgcc aggggggtgct 180
 gcgctccaag cctctg 196

<210> 20
 <211> 65
 <212> PRT
 <213> Zea mays

<400> 20
 Asn Cys Met Gly Phe Val Cys Pro Lys Tyr Phe Lys Ser Asp Leu Val
 1 5 10 15

- 11 -

Tyr Arg Thr Leu Trp Leu Gln Asp Ser Pro Thr Glu Asp Thr Thr Ser
20 25 30

Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val Arg Glu Gln Gly
35 40 45

Gly Arg Val Phe Val His Cys Cys Gln Gly Val Ser Arg Ser Thr Pro
50 55 60

Leu
65

<210> 21
<211> 2450
<212> DNA
<213> Zea mays

<400> 21
ggcgggtcct ccccgcccaa gcccggggag gggctccgcc tcgacctctc ctcgctccgg 60
tcgcaggggc gcaaggacca gatcgcttc ttcgacaagg agtgctccaa ggtcgccgac 120
cacgtctacc tcggcggcga cgccgtcgcc aagaaccgcg acatcctcag gaagaacggc 180
atcacccacg tgctcaactg tgtgggcttt gtctgcccg agtacttcaa gtcggacctt 240
gtctaccgca cctctgggt gcaggacagc ccaccgagg acatcaccag catcctgtac 300
gacgtgtttg actacttcga ggacgtcagg gagcagggcg gccgcgtgct tgtgcattgc 360
tgccaggggg tgctacgctc cacgtcgctg gtcacgcct acttgatgtg gagggaaaggc 420
cagagcttcg atgacgcctt ccagtttgtc aaggctgccc ggggatcgc aaatccaaac 480
atgggctttg catgccagct tctccagtgc cagaagcgtg tgcattgcgt tccgctgtca 540
ccaaattcag tgctcaggat gtaccgcatg gcgcctcact ccagtatgc cctctgtcat 600
ttgggtgcca aaatgtcaa tgacccatcc ccagccacc ttgactctag aggcgcgttc 660
attgtgcatg ttctctcgtc gctctatgtc tgggttgga tgaagtgtga tccgtaaatg 720
gaaaaggatg caaaggctgc tgcgtttcag gtagtgaggt atgagaagg gcaggggcac 780
atcaagggtg tgagagaagg tctggagccg caggagttct gggatgcatt ttcaagtatg 840
ccacctaatt cagatagtaa taaaagatt agcaaggacc agatcgattc agcatccaaag 900
agtgaaccag gaagccggaa aaatgagtc tatgacgtg attttgagct tgtctacaaa 960
gcaatcactg ggggagtggt cctgcattt tcaacttctg gggctggtga tgagacccat 1020
cttcagcta gagaaagtag ctggagttta ctgaggcaca agtttatctc caggctcgta 1080
gctcgtgttt attcagattc tgctctaatt aaggattttg atccacgggt acaacacctg 1140
gctgctgagg catcaacctc acctccttct ctttctccaa gtccttate atcggattca 1200
agtgtcagct cgaagtatag ttcagactca cctccttat cacctacaac tggctctcca 1260
ccatcatttg gcctctcgcc tgcttcatct aatctgacac atgctttggt gccatcatcc 1320
aggtctcccc tttctcaatc atctaataaa ggagcttcaa agccttctgg catggaatca 1380
atacactctc cttccaagac ctcttctata gcagaaagga gaggaggctt cacacttcta 1440
aagctaccat ctctccaaa ggatcttgta ttgccaccaa ggggtgccgtc tattgtattg 1500
ccaccaaggc cgccatctag tattcgagg accgaggatg cctcagataa tagtataaat 1560
ggggttaaac agctgactag tgagttttgc tcagaaaaat gcaactggtaa tagtttgagc 1620
tcgcattctg aaactagatt aattgagcgt actgacagta actcagaagt ctgcagtaat 1680
gcacaacttg tagtctacca gtggcccagc atgggaaagc taactacatt tgcacgcaag 1740
gatcttgacc cgaagtcggg ttttaatttt gttacttcca atgccatcag gagaggagaa 1800
gcagttaaaa tgggtgatgt atgggttagga ggcgaaaatg agagcagcaa gagtgttgac 1860
tccgtcgatt ggcaacaggc cactagtgat tttcttctac taaagggcct cagcaatgtt 1920
cttctgtca aggttttcaa ggagcatgaa gctgagaatc ttttggaact actgaatgtt 1980
agttaacatt aggcagtagc tatcaggata attgtagttg ctaaacaac tcaacgaagg 2040

- 12 -

catgccctcc agcatcagtc ggtaccgatg attgtcagcg aggtataaag ccacagccat 2100
 tcccttgaac ataataagct acaaacagat tccgttctgc aactgcgct catgatctat 2160
 attttgtcca gatggcagga ggctgccatg ggcgttgat cggttgcgaa ttagcactcg 2220
 tgggtgtagga gcaatcggcc gattcgggtg atattatccg ctccccgtga atgtaagctc 2280
 agatactggg agctgggtg tgcacagtta ctttttagcc taaacattct tgtacatctt 2340
 tgaaaggaac agagttgtaa tccttttgac tatgtaaag gctccattgg tcataacttc 2400
 taaacatgt ggaactcagt tgctagctgg taaaaaaaaa aaaaaaaaaa 2450

<210> 22
 <211> 661
 <212> PRT
 <213> Zea mays

<400> 22

Gly Gly Ser Ser Pro Ala Lys Pro Gly Glu Gly Leu Arg Leu Asp Leu
 1 5 10 15

Ser Ser Leu Arg Ser Gln Gly Arg Lys Asp Gln Ile Ala Phe Phe Asp
 20 25 30

Lys Glu Cys Ser Lys Val Ala Asp His Val Tyr Leu Gly Gly Asp Ala
 35 40 45

Val Ala Lys Asn Arg Asp Ile Leu Arg Lys Asn Gly Ile Thr His Val
 50 55 60

Leu Asn Cys Val Gly Phe Val Cys Pro Glu Tyr Phe Lys Ser Asp Leu
 65 70 75 80

Val Tyr Arg Thr Leu Trp Leu Gln Asp Ser Pro Thr Glu Asp Ile Thr
 85 90 95

Ser Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val Arg Glu Gln
 100 105 110

Gly Gly Arg Val Leu Val His Cys Cys Gln Gly Val Ser Arg Ser Thr
 115 120 125

Ser Leu Val Ile Ala Tyr Leu Met Trp Arg Glu Gly Gln Ser Phe Asp
 130 135 140

Asp Ala Phe Gln Phe Val Lys Ala Ala Arg Gly Ile Ala Asn Pro Asn
 145 150 155 160

Met Gly Phe Ala Cys Gln Leu Leu Gln Cys Gln Lys Arg Val His Ala
 165 170 175

Ile Pro Leu Ser Pro Asn Ser Val Leu Arg Met Tyr Arg Met Ala Pro
 180 185 190

His Ser Gln Tyr Ala Pro Leu His Leu Val Pro Lys Met Leu Asn Asp
 195 200 205

Pro Ser Pro Ala Thr Leu Asp Ser Arg Gly Ala Phe Ile Val His Val
 210 215 220

- 13 -

Leu Ser Ser Leu Tyr Val Trp Val Gly Met Lys Cys Asp Pro Val Met
 225 230 235 240
 Glu Lys Asp Ala Lys Ala Ala Ala Phe Gln Val Val Arg Tyr Glu Lys
 245 250 255
 Val Gln Gly His Ile Lys Val Val Arg Glu Gly Leu Glu Pro Gln Glu
 260 265 270
 Phe Trp Asp Ala Phe Ser Ser Met Pro Pro Asn Ser Asp Ser Asn Thr
 275 280 285
 Lys Ile Ser Lys Asp Gln Ile Asp Ser Ala Ser Lys Ser Asp Pro Gly
 290 295 300
 Ser Arg Lys Asn Glu Ser Tyr Asp Ala Asp Phe Glu Leu Val Tyr Lys
 305 310 315 320
 Ala Ile Thr Gly Gly Val Val Pro Ala Phe Ser Thr Ser Gly Ala Gly
 325 330 335
 Asp Glu Thr His Leu Pro Ala Arg Glu Ser Ser Trp Ser Leu Leu Arg
 340 345 350
 His Lys Phe Ile Ser Arg Ser Leu Ala Arg Val Tyr Ser Asp Ser Ala
 355 360 365
 Leu Met Lys Asp Phe Asp Pro Arg Val Gln His Leu Ala Ala Glu Ala
 370 375 380
 Ser Thr Ser Pro Pro Phe Leu Ser Pro Ser Ser Leu Ser Ser Asp Ser
 385 390 395 400
 Ser Val Ser Ser Lys Tyr Ser Ser Asp Ser Pro Ser Leu Ser Pro Thr
 405 410 415
 Thr Gly Ser Pro Pro Ser Phe Gly Leu Ser Pro Ala Ser Ser Asn Leu
 420 425 430
 Thr His Ala Leu Val Pro Ser Ser Arg Ser Pro Leu Ser Gln Ser Ser
 435 440 445
 Asn Glu Gly Ala Ser Lys Pro Ser Gly Met Glu Ser Ile His Ser Pro
 450 455 460
 Ser Lys Thr Ser Ser Ile Ala Glu Arg Arg Gly Gly Phe Thr Leu Leu
 465 470 475 480
 Lys Leu Pro Ser Leu Gln Lys Asp Leu Val Leu Pro Pro Arg Val Pro
 485 490 495
 Ser Ile Val Leu Pro Pro Arg Ala Pro Ser Ser Ile Arg Arg Thr Glu
 500 505 510

- 14 -

Asp Ala Ser Asp Asn Ser Thr Asn Gly Val Lys Gln Leu Thr Ser Glu
 515 520 525
 Phe Cys Ser Glu Lys Cys Thr Gly Asn Ser Leu Ser Ser His Ser Glu
 530 535 540
 Thr Arg Leu Ile Glu Arg Thr Asp Ser Asn Ser Glu Val Cys Ser Asn
 545 550 555 560
 Ala Gln Leu Val Val Tyr Gln Trp Pro Ser Met Gly Lys Leu Thr Thr
 565 570 575
 Phe Ala Arg Lys Asp Leu Asp Pro Lys Ser Val Leu Ile Phe Val Thr
 580 585 590
 Ser Asn Ala Ile Arg Arg Gly Glu Ala Val Lys Met Val Tyr Val Trp
 595 600 605
 Val Gly Gly Glu Asn Glu Ser Ser Lys Ser Val Asp Ser Val Asp Trp
 610 615 620
 Gln Gln Val Thr Ser Asp Phe Leu His Leu Lys Gly Leu Ser Asn Val
 625 630 635 640
 Leu Pro Val Lys Val Phe Lys Glu His Glu Ala Glu Asn Leu Leu Glu
 645 650 655
 Leu Leu Asn Val Ser
 660

<210> 23
 <211> 522
 <212> DNA
 <213> Lycopersicon esculentum

<400> 23
 aactgtgtgg ggtttgtatg cccagagtat ttcaagtctg atttcgtata ccggactttg 60
 tggttgcagg atagcccatc agaagatatt actagtattc tctatgatgt ttttgactac 120
 tttgaagatg tcagggagca acatgggaag gtttttggtc attgctgccca aggggtctct 180
 cggtcacact cgttggttat tgcttatcgt atgtggagag aaggacaaag ttttgatgat 240
 gcctttgagt atgtaaaggc agcaaggggt attgcggtac caaatatggg ttttgcttgt 300
 cagttattac aatgccaaaa aagggttcat gcttctcctt tgagcccaag ttcattatta 360
 aggatgtaca gaggcacc tcattcacca tacgatcctt tgcattcgt cccaaaaatg 420
 ttaaatgatc cctcaccggc agcattagat tctagaggtg catttattat acacatacct 480
 tcatcggtat atgtatggat tggtaagaaa tgtgaagcaa tc 522

<210> 24
 <211> 174
 <212> PRT
 <213> Lycopersicon esculentum

<400> 24
 Asn Cys Val Gly Phe Val Cys Pro Glu Tyr Phe Lys Ser Asp Phe Val
 1 5 10 15

- 15 -

Tyr Arg Thr Leu Trp Leu Gln Asp Ser Pro Ser Glu Asp Ile Thr Ser
 20 25 30
 Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val Arg Glu Gln His
 35 40 45
 Gly Lys Val Phe Val His Cys Cys Gln Gly Val Ser Arg Ser Thr Ser
 50 55 60
 Leu Val Ile Ala Tyr Arg Met Trp Arg Glu Gly Gln Ser Phe Asp Asp
 65 70 75 80
 Ala Phe Glu Tyr Val Lys Ala Ala Arg Gly Ile Ala Asp Pro Asn Met
 85 90 95
 Gly Phe Ala Cys Gln Leu Leu Gln Cys Gln Lys Arg Val His Ala Ser
 100 105 110
 Pro Leu Ser Pro Ser Ser Leu Leu Arg Met Tyr Arg Val Ala Pro His
 115 120 125
 Ser Pro Tyr Asp Pro Leu His Leu Val Pro Lys Met Leu Asn Asp Pro
 130 135 140
 Ser Pro Ala Ala Leu Asp Ser Arg Gly Ala Phe Ile Ile His Ile Pro
 145 150 155 160
 Ser Ser Val Tyr Val Trp Ile Gly Lys Lys Cys Glu Ala Ile
 165 170

<210> 25

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: degenerate
primer

<400> 25

gngcyttng crtcyttytc c

21

<210> 26

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: degenerate
primer

<400> 26

ytcnckngcn ggnarrtgng tytc

24



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 9/16	A3	(11) International Publication Number: WO 00/06706 (43) International Publication Date: 10 February 2000 (10.02.00)
(21) International Application Number: PCT/EP99/05413 (22) International Filing Date: 28 July 1999 (28.07.99) (30) Priority Data: 9816639.0 30 July 1998 (30.07.98) GB (71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). (72) Inventors; and (75) Inventors/Applicants (for US only): REVENKOVA, Ekaterina [RU/CH]; Peter Rot-Strasse 76, CH-4058 Basel (CH). PASZOWSKI, Jerzy [PL/CH]; Blauenweg 10, CH-4224 Nenzlingen (CH). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 4 May 2000 (04.05.00)
(54) Title: MAP KINASE PHOSPHATASE MUTANT (57) Abstract <p>The present invention relates to DNA encoding proteins contributing to the regulation of a plant's response to DNA damage. DNA according to the invention comprises an open reading frame encoding a protein characterized by a stretch of amino acids or component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO:3. Preferably the DNA encodes a MAP kinase phosphatase.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/05413

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 23039 A (CANCER RES INST ROYAL ;MARSHALL CHRISTOPHER JOHN (GB); ASHWORTH AL) 13 October 1994 (1994-10-13) figure 7D	1-12
X	& DATABASE DGENE 'Online! DERWENT Accession no. R63602, 9 May 1995 (1995-05-09) abstract	1-12
X	DATABASE SWISSPROT 'Online! Accession no. P28562, 1 December 1992 (1992-12-01) KEYES S.M. AND EMSLIE E.A.: "DUAL SPECIFICITY PROTEIN PHOSPHATASE 1 " XP002131991 abstract	1-12

-/-

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another claim or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"&" document member of the same patent family

Date of the actual completion of the international search

1 March 2000

Date of mailing of the international search report

14/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Chakravarty, A

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 99/05413

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE SWISSPROT 'Online! Accession no. P28563, 1 December 1992 (1992-12-01) CHARLES C.H. ET AL.: "DUAL SPECIFICITY PROTEIN PHOSPHATASE 1 (EC 3.1.3.48) (EC 3.1.3.16)" XP002131992 cited in the application abstract</p>	1-12
P,X	<p>DATABASE EMBL 'Online! EBI accession no. AL085836, 28 June 1999 (1999-06-28) SALANOUBAT M. ET AL.: "T7 end of BAC F11I5 of IGF library from strain Columbia of Arabidopsis thaliana" XP002131993 abstract</p>	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/05413

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9423039 A	13-10-1994	AU 677834 B	08-05-1997
		AU 6382394 A	24-10-1994
		CA 2157774 A	13-10-1994
		EP 0703984 A	03-04-1996
		JP 9501302 T	10-02-1997
		US 5958721 A	28-09-1999
		AU 696939 B	24-09-1998
		AU 1586195 A	29-08-1995
		CA 2182967 A	17-08-1995
		EP 0742827 A	20-11-1996
		WO 9521923 A	17-08-1995
		JP 9508795 T	09-09-1997

